



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

## Medical Policy

# Hematopoietic Cell Transplantation for Plasma Cell Dyscracias, Including Multiple Myeloma and POEMS Syndrome

### Table of Contents

- [Policy: Commercial](#)
- [Description](#)
- [Information Pertaining to All Policies](#)
- [Authorization Information](#)
- [Policy History](#)
- [Coding Information](#)
- [References](#)

### Policy Number: 075

BCBSA Reference Number: 8.01.17 (For Plan internal use only)

### Related Policies

- Hematopoietic Cell Transplantation for Waldenström Macroglobulinemia, #[322](#)

### Policy

#### Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity

#### MULTIPLE MYELOMA

A single or second (salvage) autologous hematopoietic cell transplantation may be **MEDICALLY NECESSARY** to treat multiple myeloma.

Tandem autologous-autologous hematopoietic cell transplantation may be **MEDICALLY NECESSARY** to treat multiple myeloma in individuals who fail to achieve at least a near-complete or very good partial response after the first transplant in the tandem sequence.

Definition of near-complete response and very good partial response

- A near complete response, as defined by the European Group for Blood and Marrow Transplant (EBMT) is the disappearance of M protein at routine electrophoresis, but positive immunofixation.
- A very good partial response has been defined as a 90% decrease in the serum paraprotein level.

Tandem transplantation with an initial round of autologous hematopoietic cell transplantation followed by a non-marrow-ablative conditioning regimen and allogeneic hematopoietic cell transplantation (i.e., reduced-intensity conditioning transplant) may be **MEDICALLY NECESSARY** to treat individuals with newly diagnosed multiple myeloma.

Allogeneic hematopoietic cell transplantation, myeloablative or nonmyeloablative, as upfront therapy of newly diagnosed multiple myeloma or as salvage therapy, is **INVESTIGATIONAL**.

#### POEMS syndrome

Autologous hematopoietic cell transplantation may be considered **MEDICALLY NECESSARY** to treat disseminated POEMS syndrome.\*

\*Patients with disseminated POEMS syndrome may have diffuse sclerotic lesions or disseminated bone marrow involvement.

Allogeneic and tandem hematopoietic cell transplantation are **INVESTIGATIONAL** to treat POEMS syndrome.

## Prior Authorization Information

### Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

### Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	Prior authorization is <b>required</b> .
Commercial PPO and Indemnity	Prior authorization is <b>required</b> .

### Requesting Prior Authorization Using Authorization Manager

Providers will need to use [Authorization Manager](#) to submit initial authorization requests for services. Authorization Manager, available 24/7, is the quickest way to review authorization requirements, request authorizations, submit clinical documentation, check existing case status, and view/print the decision letter. For commercial members, the requests must meet medical policy guidelines.

To ensure the service request is processed accurately and quickly:

- Enter the facility's NPI or provider ID for where services are being performed.
- Enter the appropriate surgeon's NPI or provider ID as the servicing provider, *not* the billing group.

### Authorization Manager Resources

Refer to our [Authorization Manager](#) page for tips, guides, and video demonstrations.

## CPT Codes / HCPCS Codes / ICD Codes

*Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.*

*Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.*

*The following codes are included below for informational purposes only; this is not an all-inclusive list.*

**The above medical necessity criteria MUST be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:**

### CPT Codes

CPT codes:	Code Description
38241	Bone marrow or blood-derived peripheral stem-cell transplantation; autologous

### HCPCS Codes

<b>HCPCS codes:</b>	<b>Code Description</b>
S2150	Bone marrow or blood-derived peripheral stem-cell (peripheral or umbilical), allogeneic or autologous, harvesting, transplantation, and related complications including pheresis and cell preparation/storage; marrow ablative therapy; drugs, supplies, hospitalization with outpatient follow-up; medical/surgical, diagnostic, emergency, and rehabilitative services; and the number of days of pre- and post-transplant care in the global definition

### **ICD-10-PCS Procedure Codes**

<b>ICD-10-PCS procedure codes:</b>	<b>Code Description</b>
30233G0	Transfusion of Autologous Bone Marrow into Peripheral Vein, Percutaneous Approach
30233X0	Transfusion of Autologous Cord Blood Stem Cells into Peripheral Vein, Percutaneous Approach
30233Y0	Transfusion of Autologous Hematopoietic Stem Cells into Peripheral Vein, Percutaneous Approach
30243G0	Transfusion of Autologous Bone Marrow into Central Vein, Percutaneous Approach
30243X0	Transfusion of Autologous Cord Blood Stem Cells into Central Vein, Percutaneous Approach
30243Y0	Transfusion of Autologous Hematopoietic Stem Cells into Central Vein, Percutaneous Approach
30263G0	Transfusion of Autologous Bone Marrow into Central Artery, Percutaneous Approach
30263X0	Transfusion of Autologous Cord Blood Stem Cells into Central Artery, Percutaneous Approach
30263Y0	Transfusion of Autologous Hematopoietic Stem Cells into Central Artery, Percutaneous Approach
3E04305	Introduction of Other Antineoplastic into Central Vein, Percutaneous Approach

**The following CPT, HCPCS and ICD Procedure codes are considered investigational for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:**

### **CPT Codes**

<b>CPT codes:</b>	<b>Code Description</b>
38240	Bone marrow or blood-derived peripheral stem-cell transplantation; allogeneic

### **HCPCS Codes**

<b>HCPCS codes:</b>	<b>Code Description</b>
S2142	Cord blood-derived stem-cell transplantation, allogeneic

### **ICD-10-PCS Procedure Codes**

<b>ICD-10-PCS procedure codes:</b>	<b>Code Description</b>
30233G1	Transfusion of Nonautologous Bone Marrow into Peripheral Vein, Percutaneous Approach
30233X1	Transfusion of Nonautologous Cord Blood Stem Cells into Peripheral Vein, Percutaneous Approach

30233Y1	Transfusion of Nonautologous Hematopoietic Stem Cells into Peripheral Vein, Percutaneous Approach
30243G1	Transfusion of Nonautologous Bone Marrow into Central Vein, Percutaneous Approach
30243X1	Transfusion of Nonautologous Cord Blood Stem Cells into Central Vein, Percutaneous Approach
30243Y1	Transfusion of Nonautologous Hematopoietic Stem Cells into Central Vein, Percutaneous Approach
30263G1	Transfusion of Nonautologous Bone Marrow into Central Artery, Percutaneous Approach
30263X1	Transfusion of Nonautologous Cord Blood Stem Cells into Central Artery, Percutaneous Approach
30263Y1	Transfusion of Nonautologous Hematopoietic Stem Cells into Central Artery, Percutaneous Approach

## Description

### Multiple Myeloma

Multiple myeloma (MM) is a systemic malignancy of plasma cells that represents approximately 18% of all hematologic cancers in the United States. It is treatable but rarely curable. At diagnosis, most patients have generalized disease, and the selection of treatment is influenced by patient age, general health, prior therapy, and the presence of disease complications.<sup>1,2,3,4</sup>

The disease is staged by estimating tumor mass, based on various clinical parameters such as hemoglobin, serum calcium, number of lytic bone lesions, and the presence or absence of renal failure. Multiple myeloma usually evolves from an asymptomatic premalignant stage (termed *monoclonal gammopathy of undetermined significance*). Treatment is usually reserved for patients with symptomatic disease (usually progressive myeloma), whereas asymptomatic patients are observed because there is little evidence that early treatment of asymptomatic MM prolongs survival compared with therapy delivered at the time of symptoms or end-organ damage.<sup>1,2</sup> In some patients, an intermediate asymptomatic but more advanced premalignant stage is recognized and referred to as smoldering MM. The overall risk of disease progression from smoldering to symptomatic MM is 10% per year for the first 5 years, approximately 3% per year for the next 5 years, and 1% for the next 10 years.<sup>1,2</sup>

### Polyneuropathy, Organomegaly, Endocrinopathy, Monoclonal gammopathy, and Skin abnormalities (POEMS) Syndrome

POEMS syndrome (also known as osteosclerotic myeloma, Crow-Fukase syndrome, or Takatsuki syndrome) is a rare, paraneoplastic disorder secondary to a plasma cell dyscrasia.<sup>5,6</sup> This complex, multiorgan disease was first described in 1938, but the acronym POEMS was coined in 1980, reflecting hallmark characteristics of the syndrome: polyneuropathy, organomegaly, endocrinopathy, M protein, and skin changes.<sup>7</sup> No single test establishes the presence of POEMS syndrome. Its pathogenesis is undefined, although some evidence has suggested it is mediated by an imbalance of proinflammatory cytokines including interleukin (IL)-1 $\beta$ , IL-6, and tumor necrosis factor  $\alpha$ ; vascular endothelial growth factor may also be involved.<sup>6,8</sup> However, specific criteria have been established, and the syndrome may entail other findings in the constellation of signs and symptoms, as shown in Table 1.<sup>9</sup> Both mandatory major criteria, at least 1 of the other major criteria, and at least 1 of the minor criteria are necessary for diagnosis.

**Table 1. Criteria and Associations for POEMS Syndrome**

Mandatory Major Criteria	Other Major Criteria	Minor Criteria	Other Symptoms and Signs
Polyneuropathy	Castleman disease	Organomegaly (splenomegaly, hepatomegaly, lymphadenopathy)	Pulmonary hypertension/restrictive lung disease

Monoclonal plasma-proliferative disorder	Sclerotic bone lesions	Extravascular volume overload (edema, pleural effusion, ascites)	Clubbing
	Vascular endothelial growth factor elevation	Endocrinopathy (adrenal, thyroid, pituitary, gonadal, parathyroid, pancreatic)	Thrombotic diatheses
		Skin changes (hyperpigmentation, hypertrichosis, plethora, hemangiomas, white nails)	Weight loss
		Papilledema	Low vitamin B <sub>12</sub> levels
		Thrombocytosis/polycythemia	Diarrhea
			Hyperhidrosis

The prevalence of POEMS syndrome is unclear. A national survey in Japan showed a prevalence of about 0.3 per 100,000.<sup>10</sup> Other large series have been described in the United States, France, China, and India.<sup>9</sup> In general, patients with POEMS have superior overall survival (OS) compared with that of MM (nearly 14 years in a large series).<sup>8</sup> However, given the rarity of POEMS, there is a paucity of randomized controlled trial (RCT) evidence for POEMS therapies.<sup>9</sup> Numerous approaches have been tried, including ionizing radiation, plasmapheresis, intravenous immunoglobulin, interferon- $\alpha$ , corticosteroids, alkylating agents, tamoxifen, trans-retinoic acid, and high-dose chemotherapy with autologous hematopoietic cell transplantation (HCT) support. Optimal treatment involves eliminating the plasma cell clone (eg, by surgical excision or local radiotherapy for an isolated plasmacytoma) or systemic chemotherapy in patients with disseminated disease (eg, medullary disease or multiple plasmacytomas). Given the underlying plasma cell dyscrasia of POEMS syndrome, newer approaches to MM, including bortezomib, lenalidomide, and thalidomide, have also been investigated.

**Hematopoietic Cell Transplantation**

Hematopoietic cell transplantation is a procedure in which hematopoietic stem cells are intravenously infused to restore bone marrow and immune function in cancer patients who receive bone marrow-toxic doses of cytotoxic drugs with or without whole-body radiotherapy. Hematopoietic stem cells may be obtained from the transplant recipient (autologous HCT) or a donor (allogeneic HCT [allo-HCT]). They can be harvested from bone marrow, peripheral blood, or umbilical cord blood shortly after delivery of neonates.

Immunologic compatibility between infused hematopoietic stem cells and the recipient is not an issue in autologous HCT. In allo-HCT, immunologic compatibility between donor and patient is a critical factor for achieving a successful outcome. Compatibility is established by typing of human leukocyte antigens (HLA) using cellular, serologic, or molecular techniques. HLA refers to the gene complex expressed at the HLA-A, -B, and -DR (antigen-D related) loci on each arm of chromosome 6. An acceptable donor will match the patient at all or most of the HLA loci.

**Conditioning for Hematopoietic Cell Transplantation**

**Conventional Conditioning**

The conventional (“classical”) practice of allo-HCT involves administration of cytotoxic agents (e.g., cyclophosphamide, busulfan) with or without total body irradiation at doses sufficient to cause bone marrow ablation in the recipient. The beneficial treatment effect of this procedure is due to a combination of the initial eradication of malignant cells and subsequent graft-versus-malignancy effect mediated by non-self-immunologic effector cells. While the slower graft-versus-malignancy effect is considered the potentially curative component, it may be overwhelmed by existing disease in the absence of pretransplant conditioning. Intense conditioning regimens are limited to patients who are sufficiently medically fit to tolerate substantial adverse effects. These include opportunistic infections secondary to loss of endogenous bone marrow function and organ damage or failure caused by cytotoxic drugs.

Subsequent to graft infusion in allo-HCT, immunosuppressant drugs are required to minimize graft rejection and graft-versus-host disease, which increases susceptibility to opportunistic infections.

The success of autologous HCT is predicated on the potential of cytotoxic chemotherapy, with or without radiotherapy, to eradicate cancerous cells from the blood and bone marrow. This permits subsequent engraftment and repopulation of the bone marrow with presumably normal hematopoietic stem cells obtained from the patient before undergoing bone marrow ablation. Therefore, autologous HCT is typically performed as consolidation therapy when the patient's disease is in complete remission. Patients who undergo autologous HCT are also susceptible to chemotherapy-related toxicities and opportunistic infections before engraftment, but not graft-versus-host disease.

### **Reduced-Intensity Conditioning Allogeneic Hematopoietic Cell Transplantation**

Reduced-intensity conditioning (RIC) refers to the pretransplant use of lower doses of cytotoxic drugs or less intense regimens of radiotherapy than are used in traditional full-dose myeloablative conditioning treatments. Although the definition of RIC is variable, with numerous versions employed, all regimens seek to balance the competing effects of relapse due to residual disease and non-relapse mortality. The goal of RIC is to reduce disease burden and to minimize associated treatment-related morbidity and non-relapse mortality in the period during which the beneficial graft-versus-malignancy effect of allogeneic transplantation develops. Reduced-intensity conditioning regimens range from nearly total myeloablative to minimally myeloablative with lymphoablation, with intensity tailored to specific diseases and patient condition. Patients who undergo RIC with allo-HCT initially demonstrate donor cell engraftment and bone marrow mixed chimerism. Most will subsequently convert to full-donor chimerism. In this review, the term RIC will refer to all conditioning regimens intended to be nonmyeloablative.

### **Multiple Myeloma Treatment Overview**

In the prechemotherapy era, the median survival for a patient diagnosed with MM was approximately 7 months. After the introduction of chemotherapy (eg, the alkylating agent melphalan in the 1960s), prognosis improved, with a median survival of 24 to 30 months and 10-year survival of 3%. In a large group of patients with newly diagnosed MM, there was no difference in OS reported during a 24-year period from 1971 to 1994, with a trend toward improvement from 1995 to 2000, and a statistically significant benefit in OS from 2001 to 2006.<sup>2</sup> These data suggested that autologous HCT was responsible for the trends from 1994 to 2000, while novel agents have contributed to the improvement since 2001.

The introduction of novel agents and better prognostic indicators has been the major advances in the treatment of this disease.<sup>11</sup> Novel agents such as the proteasome inhibitors (eg, bortezomib), the monoclonal antibody daratumumab, and the immunomodulatory derivatives thalidomide and lenalidomide first showed efficacy in relapsed and refractory myeloma and now have been integrated into first-line regimens.<sup>11,12,13</sup> With the introduction of these novel treatments, it is now expected that most patients with MM will respond to initial therapy, and only a small minority will have refractory disease.<sup>14</sup>

## **Summary**

### **Description**

Multiple myeloma (MM) is a systemic malignancy of plasma cells that represents approximately 10% of all hematologic cancers. POEMS syndrome, characterized by polyneuropathy, organomegaly, endocrinopathy, M protein, and skin changes, is a rare, paraneoplastic disorder secondary to a plasma cell dyscrasia. Plasma cell dyscrasias are treatable but rarely curable. In some cases, autologous or allogeneic hematopoietic cell transplantation (HCT) is considered as therapy.

### **Summary of Evidence**

#### **Newly Diagnosed Multiple Myeloma**

For individuals who have newly diagnosed multiple myeloma (MM) who receive autologous hematopoietic cell transplantation (HCT) as initial treatment, the evidence includes reviews, a retrospective study, and several prospective randomized controlled trials (RCTs) that compare high-dose chemotherapy plus autologous HCT to standard chemotherapy regimens or regimens containing newer MM agents. Relevant

outcomes are overall survival (OS) and treatment-related morbidity. In general, the evidence has suggested OS rates are improved with autologous HCT compared with conventional chemotherapy in this setting. Limitations of the published evidence include patient heterogeneity, variability in treatment protocols, short follow-up periods, inconsistency in reporting important health outcomes, and inconsistency in reporting or collecting outcomes. Recent RCTs comparing high-dose chemotherapy plus autologous HCT to regimens that include novel MM agents have also shown that high-dose chemotherapy plus autologous HCT improves progression-free survival (PFS). The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have newly diagnosed MM who receive tandem autologous HCT, the evidence includes several RCTs. Relevant outcomes are OS and treatment-related morbidity. Compared with single autologous HCT, RCTs have generally found that tandem autologous HCT improves OS and recurrence-free survival in newly diagnosed MM. Two recent RCTs found conflicting results on the benefit of tandem autologous HCT versus single autologous HCT; however, the study that found no additional benefit with tandem autologous HCT had a higher rate of nonadherence to the second planned HCT. Differences in initial therapy regimens between trials may also have led to conflicting results. Several RCTs and one retrospective study compared reduced-intensity conditioning (RIC) allogeneic HCT (allo-HCT) following a first autologous HCT with single or tandem autologous transplants. The RCTs were based on genetic randomization (ie, patients with a human leukocyte antigen-identical sibling were offered RIC allo-HCT following autologous HCT, whereas other patients underwent either 1 or 2 autologous transplants). Although the body of evidence has shown inconsistencies regarding OS and disease-free survival rates, some studies have shown a survival benefit with tandem autologous HCT followed by RIC allo-HCT, although at the cost of higher transplant-related mortality compared with conventional treatments. Factors across studies that may account for differing trial results include different study designs, nonuniform preparative regimens, different patient characteristics (including risk stratification), and criteria for advancing to a second transplant. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have newly diagnosed MM who receive allo-HCT as initial or salvage treatment, the evidence includes nonrandomized studies. Relevant outcomes are OS and treatment-related morbidity. Studies have reported on patients with both myeloablative conditioning and RIC. Limitations of the published evidence include patient sample heterogeneity, variability in treatment protocols, short follow-up periods, inconsistency in reporting important health outcomes, and inconsistency in reporting or collecting outcomes. Nonmyeloablative allo-HCT as first-line therapy is associated with lower transplant-related mortality but a greater risk of relapse; convincing evidence is lacking that allo-HCT improves survival better than autologous HCT. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

### **Relapsed or Refractory Multiple Myeloma**

For individuals who have relapsed MM after failing an autologous HCT who receive autologous HCT, the evidence includes RCTs, retrospective studies, and reviews summarizing recent studies on a second autologous HCT in relapsed myeloma. Relevant outcomes are OS and treatment-related morbidity. Despite some limitations of the published evidence, including patient sample heterogeneity, variability in treatment protocols, and short follow-up periods, the available trial evidence has suggested OS rates are improved with autologous HCT compared with conventional chemotherapy or continuous lenalidomide plus dexamethasone in this setting. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have refractory multiple myeloma after failing a first HCT who receive tandem autologous HCT, the evidence includes systematic reviews and a retrospective study. Relevant outcomes are OS and treatment-related morbidity. The evidence has shown tandem autologous HCT improves OS rates in this setting. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

## **Polyneuropathy, Organomegaly, Endocrinopathy, Monoclonal gammopathy, and Skin abnormalities (POEMS) Syndrome**

For individuals who have POEMS syndrome who receive HCT, the evidence includes retrospective cohort studies, case reports, and case series. Relevant outcomes are OS and treatment-related morbidity. No RCTs of HCT of any type have been performed in patients with POEMS syndrome of any severity, nor is it likely such studies will be performed because of the rarity of this condition. Available case reports and series are subject to selection bias and are heterogeneous concerning treatment approaches and peritransplant support. However, for patients with disseminated POEMS syndrome, a chain of evidence and contextual factors related to the disease and MM would suggest improvement in health outcomes with autologous HCT. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

### **Policy History**

<b>Date</b>	<b>Action</b>
3/2024	Annual policy review. References updated. Policy statements unchanged.
9/2023	Policy clarified to include prior authorization requests using Authorization Manager.
3/2023	Annual policy review. Minor editorial refinements to policy statements; intent unchanged.
2/2022	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
3/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
1/2021	Medicare information removed. See MP #132 Medicare Advantage Management for local coverage determination and national coverage determination reference.
4/2020	Bone marrow harvesting codes were removed. Outpatient prior authorization is not required.
3/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2019	Clarified coding information
3/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
1/2019	Outpatient prior authorization is required for all commercial products. Effective 1/1/2019.
2/2018	Annual policy review. New references added. Title clarified.
1/2018	Clarified coding information.
11/2015	Annual policy review. New references added
4/2015	Annual policy review. New references added
1/2015	Clarified coding information.
5/2014	Updated Coding section with ICD10 procedure and diagnosis codes, effective 10/2015.
2/2014	Annual policy review. New medically necessary and investigational indications described; policy title changed. Effective 2/1/2014.
12/2012	Updated to add new CPT code 38243.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
12/2011	Minor change to policy statements (added phrase “in the tandem sequence” to the medically necessary tandem autologous-autologous statement).
7/2011	Medical Policy Group – Hematology and Oncology. No changes to policy statements.
9/2010	Medical Policy Group – Hematology and Oncology. No changes to policy statements.
9/1/2010	Medical Policy 075 effective 9/1/2010.



## Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

## References

1. Kyle RA, Rajkumar SV. Multiple myeloma. *Blood*. Mar 15 2008; 111(6): 2962-72. PMID 18332230
2. Palumbo A, Rajkumar SV. Treatment of newly diagnosed myeloma. *Leukemia*. Mar 2009; 23(3): 449-56. PMID 19005483
3. Durie BG, Harousseau JL, Miguel JS, et al. International uniform response criteria for multiple myeloma. *Leukemia*. Sep 2006; 20(9): 1467-73. PMID 16855634
4. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Multiple Myeloma. Version 2.2024.  
[https://www.nccn.org/professionals/physician\\_gls/pdf/myeloma.pdf](https://www.nccn.org/professionals/physician_gls/pdf/myeloma.pdf). Accessed November 27, 2023.
5. Dispenzieri A. Long-term outcomes after autologous stem cell transplantation in patients with POEMS syndrome. *Clin Adv Hematol Oncol*. Nov 2012; 10(11): 744-6. PMID 23271262
6. Dispenzieri A. POEMS syndrome: update on diagnosis, risk-stratification, and management. *Am J Hematol*. Aug 2012; 87(8): 804-14. PMID 22806697
7. Bardwick PA, Zvaifler NJ, Gill GN, et al. Plasma cell dyscrasia with polyneuropathy, organomegaly, endocrinopathy, M protein, and skin changes: the POEMS syndrome. Report on two cases and a review of the literature. *Medicine (Baltimore)*. Jul 1980; 59(4): 311-22. PMID 6248720
8. Dispenzieri A, Kyle RA, Lacy MQ, et al. POEMS syndrome: definitions and long-term outcome. *Blood*. Apr 01 2003; 101(7): 2496-506. PMID 12456500
9. Dispenzieri A. POEMS Syndrome: 2019 Update on diagnosis, risk-stratification, and management. *Am J Hematol*. Jul 2019; 94(7): 812-827. PMID 31012139
10. Nasu S, Misawa S, Sekiguchi Y, et al. Different neurological and physiological profiles in POEMS syndrome and chronic inflammatory demyelinating polyneuropathy. *J Neurol Neurosurg Psychiatry*. May 2012; 83(5): 476-9. PMID 22338030
11. Reece DE. Recent trends in the management of newly diagnosed multiple myeloma. *Curr Opin Hematol*. Jul 2009; 16(4): 306-12. PMID 19491669
12. Qiao SK, Guo XN, Ren JH, et al. Efficacy and Safety of Lenalidomide in the Treatment of Multiple Myeloma: A Systematic Review and Meta-analysis of Randomized Controlled Trials. *Chin Med J (Engl)*. May 05 2015; 128(9): 1215-22. PMID 25947406
13. Rajkumar SV, Kumar S. Multiple myeloma current treatment algorithms. *Blood Cancer J*. Sep 28 2020; 10(9): 94. PMID 32989217
14. Fonseca R. Strategies for risk-adapted therapy in myeloma. *Hematology Am Soc Hematol Educ Program*. 2007: 304-10. PMID 18024644
15. Rajkumar SV. Multiple myeloma: 2020 update on diagnosis, risk-stratification and management. *Am J Hematol*. May 2020; 95(5): 548-567. PMID 32212178
16. Mian H, Mian OS, Rochweg B, et al. Autologous stem cell transplant in older patients (age  $\geq$  65) with newly diagnosed multiple myeloma: A systematic review and meta-analysis. *J Geriatr Oncol*. Jan 2020; 11(1): 93-99. PMID 31153809
17. Koreth J, Cutler CS, Djulbegovic B, et al. High-dose therapy with single autologous transplantation versus chemotherapy for newly diagnosed multiple myeloma: A systematic review and meta-analysis of randomized controlled trials. *Biol Blood Marrow Transplant*. Feb 2007; 13(2): 183-96. PMID 17241924
18. Richardson PG, Jacobus SJ, Weller EA, et al. Triplet Therapy, Transplantation, and Maintenance until Progression in Myeloma. *N Engl J Med*. Jul 14 2022; 387(2): 132-147. PMID 35660812
19. Cavo M, Gay F, Beksac M, et al. Autologous haematopoietic stem-cell transplantation versus bortezomib-melphalan-prednisone, with or without bortezomib-lenalidomide-dexamethasone consolidation therapy, and lenalidomide maintenance for newly diagnosed multiple myeloma

- (EMN02/HO95): a multicentre, randomised, open-label, phase 3 study. *Lancet Haematol.* Jun 2020; 7(6): e456-e468. PMID 32359506
20. Attal M, Lauwers-Cances V, Hulin C, et al. Lenalidomide, Bortezomib, and Dexamethasone with Transplantation for Myeloma. *N Engl J Med.* Apr 06 2017; 376(14): 1311-1320. PMID 28379796
  21. Gay F, Oliva S, Petrucci MT, et al. Chemotherapy plus lenalidomide versus autologous transplantation, followed by lenalidomide plus prednisone versus lenalidomide maintenance, in patients with multiple myeloma: a randomised, multicentre, phase 3 trial. *Lancet Oncol.* Dec 2015; 16(16): 1617-29. PMID 26596670
  22. Attal M, Harousseau JL. The role of high-dose therapy with autologous stem cell support in the era of novel agents. *Semin Hematol.* Apr 2009; 46(2): 127-32. PMID 19389496
  23. Attal M, Harousseau JL, Stoppa AM, et al. A prospective, randomized trial of autologous bone marrow transplantation and chemotherapy in multiple myeloma. *Intergroupe Français du Myélome. N Engl J Med.* Jul 11 1996; 335(2): 91-7. PMID 8649495
  24. Barlogie B, Kyle RA, Anderson KC, et al. Standard chemotherapy compared with high-dose chemoradiotherapy for multiple myeloma: final results of phase III US Intergroup Trial S9321. *J Clin Oncol.* Feb 20 2006; 24(6): 929-36. PMID 16432076
  25. Bladé J, Rosiñol L, Sureda A, et al. High-dose therapy intensification compared with continued standard chemotherapy in multiple myeloma patients responding to the initial chemotherapy: long-term results from a prospective randomized trial from the Spanish cooperative group PETHEMA. *Blood.* Dec 01 2005; 106(12): 3755-9. PMID 16105975
  26. Child JA, Morgan GJ, Davies FE, et al. High-dose chemotherapy with hematopoietic stem-cell rescue for multiple myeloma. *N Engl J Med.* May 08 2003; 348(19): 1875-83. PMID 12736280
  27. Femand JP, Ravaud P, Chevret S, et al. High-dose therapy and autologous peripheral blood stem cell transplantation in multiple myeloma: up-front or rescue treatment? Results of a multicenter sequential randomized clinical trial. *Blood.* Nov 01 1998; 92(9): 3131-6. PMID 9787148
  28. Palumbo A, Brinthen S, Petrucci MT, et al. Intermediate-dose melphalan improves survival of myeloma patients aged 50 to 70: results of a randomized controlled trial. *Blood.* Nov 15 2004; 104(10): 3052-7. PMID 15265788
  29. Marini C, Maia T, Bergantim R, et al. Real-life data on safety and efficacy of autologous stem cell transplantation in elderly patients with multiple myeloma. *Ann Hematol.* Feb 2019; 98(2): 369-379. PMID 30368589
  30. Attal M, Harousseau JL, Facon T, et al. Single versus double autologous stem-cell transplantation for multiple myeloma. *N Engl J Med.* Dec 25 2003; 349(26): 2495-502. PMID 14695409
  31. Stadtmauer EA. Multiple myeloma, 2004--one or two transplants?. *N Engl J Med.* Dec 25 2003; 349(26): 2551-3. PMID 14695416
  32. Cavo M, Tosi P, Zamagni E, et al. Prospective, randomized study of single compared with double autologous stem-cell transplantation for multiple myeloma: Bologna 96 clinical study. *J Clin Oncol.* Jun 10 2007; 25(17): 2434-41. PMID 17485707
  33. Stadtmauer EA, Pasquini MC, Blackwell B, et al. Autologous Transplantation, Consolidation, and Maintenance Therapy in Multiple Myeloma: Results of the BMT CTN 0702 Trial. *J Clin Oncol.* Mar 01 2019; 37(7): 589-597. PMID 30653422
  34. Continued, Long-Term Follow-Up and Lenalidomide Maintenance Therapy for Patients on BMT CTN 0702 Protocol (BMT CTN 07LT). *Clinicaltrials.gov.* Updated May 11, 2020. <https://clinicaltrials.gov/ct2/show/NCT02322320>. Accessed November 28, 2023.
  35. Villalba A, Gonzalez-Rodriguez AP, Arzuaga-Mendez J, et al. Single versus tandem autologous stem-cell transplantation in patients with newly diagnosed multiple myeloma and high-risk cytogenetics. A retrospective, open-label study of the PETHEMA/Spanish Myeloma Group (GEM). *Leuk Lymphoma.* Dec 2022; 63(14): 3438-3447. PMID 36124538
  36. Garban F, Attal M, Michallet M, et al. Prospective comparison of autologous stem cell transplantation followed by dose-reduced allograft (IFM99-03 trial) with tandem autologous stem cell transplantation (IFM99-04 trial) in high-risk de novo multiple myeloma. *Blood.* May 01 2006; 107(9): 3474-80. PMID 16397129
  37. Moreau P, Garban F, Attal M, et al. Long-term follow-up results of IFM99-03 and IFM99-04 trials comparing nonmyeloablative allotransplantation with autologous transplantation in high-risk de novo multiple myeloma. *Blood.* Nov 01 2008; 112(9): 3914-5. PMID 18948589

38. Bruno B, Rotta M, Patriarca F, et al. A comparison of allografting with autografting for newly diagnosed myeloma. *N Engl J Med*. Mar 15 2007; 356(11): 1110-20. PMID 17360989
39. Rosiñol L, Pérez-Simón JA, Sureda A, et al. A prospective PETHEMA study of tandem autologous transplantation versus autograft followed by reduced-intensity conditioning allogeneic transplantation in newly diagnosed multiple myeloma. *Blood*. Nov 01 2008; 112(9): 3591-3. PMID 18612103
40. Björkstrand B, Iacobelli S, Hegenbart U, et al. Autologous stem cell transplantation (ASCT) versus ASCT followed by reduced-intensity conditioning allogeneic SCT with identical sibling donor in previously untreated multiple myeloma: preliminary analysis of a prospective controlled trial by the EBMT [abstract]. *Bone Marrow Transplant*. 2008;41:S38.
41. Gahrton G, Iacobelli S, Björkstrand B, et al. Autologous/reduced-intensity allogeneic stem cell transplantation vs autologous transplantation in multiple myeloma: long-term results of the EBMT-NMAM2000 study. *Blood*. Jun 20 2013; 121(25): 5055-63. PMID 23482933
42. Krishnan A, Pasquini MC, Logan B, et al. Autologous haemopoietic stem-cell transplantation followed by allogeneic or autologous haemopoietic stem-cell transplantation in patients with multiple myeloma (BMT CTN 0102): a phase 3 biological assignment trial. *Lancet Oncol*. Dec 2011; 12(13): 1195-203. PMID 21962393
43. Maffini E, Storer BE, Sandmaier BM, et al. Long-term follow up of tandem autologous-allogeneic hematopoietic cell transplantation for multiple myeloma. *Haematologica*. Feb 2019; 104(2): 380-391. PMID 30262560
44. Giral S, Koehne G. Allogeneic hematopoietic stem cell transplantation for multiple myeloma: what place, if any?. *Curr Hematol Malig Rep*. Dec 2013; 8(4): 284-90. PMID 24146203
45. Giral S, Costa L, Schriber J, et al. Optimizing autologous stem cell mobilization strategies to improve patient outcomes: consensus guidelines and recommendations. *Biol Blood Marrow Transplant*. Mar 2014; 20(3): 295-308. PMID 24141007
46. Harsouseau JL. The allogeneic dilemma. *Bone Marrow Transplant*. Dec 2007; 40(12): 1123-8. PMID 17680016
47. Crawley C, Iacobelli S, Björkstrand B, et al. Reduced-intensity conditioning for myeloma: lower nonrelapse mortality but higher relapse rates compared with myeloablative conditioning. *Blood*. Apr 15 2007; 109(8): 3588-94. PMID 17158231
48. Gahrton G, Björkstrand B. Allogeneic transplantation in multiple myeloma. *Haematologica*. Sep 2008; 93(9): 1295-300. PMID 18757850
49. Ziogas DC, Terpos E, Dimopoulos MA. When to recommend a second autograft in patients with relapsed myeloma?. *Leuk Lymphoma*. Apr 2017; 58(4): 781-787. PMID 27894207
50. Garderet L, Cook G, Auner HW, et al. Treatment options for relapse after autograft in multiple myeloma - report from an EBMT educational meeting. *Leuk Lymphoma*. Apr 2017; 58(4): 797-808. PMID 27650125
51. Goldschmidt H, Baertsch MA, Schlenzka J, et al. Salvage autologous transplant and lenalidomide maintenance vs. lenalidomide/dexamethasone for relapsed multiple myeloma: the randomized GMMG phase III trial ReLApsE. *Leukemia*. Apr 2021; 35(4): 1134-1144. PMID 32694619
52. Cook G, Williams C, Brown JM, et al. High-dose chemotherapy plus autologous stem-cell transplantation as consolidation therapy in patients with relapsed multiple myeloma after previous autologous stem-cell transplantation (NCRI Myeloma X Relapse [Intensive trial]): a randomised, open-label, phase 3 trial. *Lancet Oncol*. Jul 2014; 15(8): 874-85. PMID 24948586
53. Cook G, Ashcroft AJ, Cairns DA, et al. The effect of salvage autologous stem-cell transplantation on overall survival in patients with relapsed multiple myeloma (final results from BSBMT/UKMF Myeloma X Relapse [Intensive]): a randomised, open-label, phase 3 trial. *Lancet Haematol*. Jul 2016; 3(7): e340-51. PMID 27374467
54. Ikeda T, Mori K, Kawamura K, et al. Comparison between autologous and allogeneic stem cell transplantation as salvage therapy for multiple myeloma relapsing/progressing after autologous stem cell transplantation. *Hematol Oncol*. Dec 2019; 37(5): 586-594. PMID 31674032
55. Michaelis LC, Saad A, Zhong X, et al. Salvage second hematopoietic cell transplantation in myeloma. *Biol Blood Marrow Transplant*. May 2013; 19(5): 760-6. PMID 23298856
56. Qazilbash MH, Saliba R, De Lima M, et al. Second autologous or allogeneic transplantation after the failure of first autograft in patients with multiple myeloma. *Cancer*. Mar 01 2006; 106(5): 1084-9. PMID 16456814

57. Hahn T, Wingard JR, Anderson KC, et al. The role of cytotoxic therapy with hematopoietic stem cell transplantation in the therapy of multiple myeloma: an evidence-based review. *Biol Blood Marrow Transplant.* Jan 2003; 9(1): 4-37. PMID 12533739
58. McCarthy PL, Holstein SA. Role of stem cell transplant and maintenance therapy in plasma cell disorders. *Hematology Am Soc Hematol Educ Program.* Dec 02 2016; 2016(1): 504-511. PMID 27913522
59. Olin RL, Vogl DT, Porter DL, et al. Second auto-SCT is safe and effective salvage therapy for relapsed multiple myeloma. *Bone Marrow Transplant.* Mar 2009; 43(5): 417-22. PMID 18850013
60. Kuwabara S, Dispenzieri A, Arimura K, et al. Treatment for POEMS (polyneuropathy, organomegaly, endocrinopathy, M-protein, and skin changes) syndrome. *Cochrane Database Syst Rev.* Jun 13 2012; 2012(6): CD006828. PMID 22696361
61. Autore F, Innocenti I, Luigetti M, et al. Autologous peripheral blood stem cell transplantation and the role of lenalidomide in patients affected by poems syndrome. *Hematol Oncol.* Apr 2018; 36(2): 392-398. PMID 28913957
62. Cook G, Iacobelli S, van Biezen A, et al. High-dose therapy and autologous stem cell transplantation in patients with POEMS syndrome: a retrospective study of the Plasma Cell Disorder sub-committee of the Chronic Malignancy Working Party of the European Society for Blood Marrow Transplantation. *Haematologica.* Jan 2017; 102(1): 160-167. PMID 27634201
63. D'Souza A, Lacy M, Gertz M, et al. Long-term outcomes after autologous stem cell transplantation for patients with POEMS syndrome (osteosclerotic myeloma): a single-center experience. *Blood.* Jul 05 2012; 120(1): 56-62. PMID 22611150
64. Jang IY, Yoon DH, Kim S, et al. Advanced POEMS syndrome treated with high-dose melphalan followed by autologous blood stem cell transplantation: a single-center experience. *Blood Res.* Mar 2014; 49(1): 42-8. PMID 24724066
65. Mikhael J, Ismaila N, Cheung MC, et al. Treatment of Multiple Myeloma: ASCO and CCO Joint Clinical Practice Guideline. *J Clin Oncol.* May 10 2019; 37(14): 1228-1263. PMID 30932732
66. Shah N, Callander N, Ganguly S, et al. Hematopoietic Stem Cell Transplantation for Multiple Myeloma: Guidelines from the American Society for Blood and Marrow Transplantation. *Biol Blood Marrow Transplant.* Jul 2015; 21(7): 1155-66. PMID 25769794
67. Dhakal B, Shah N, Kansagra A, et al. ASTCT Clinical Practice Recommendations for Transplantation and Cellular Therapies in Multiple Myeloma. *Transplant Cell Ther.* Jun 2022; 28(6): 284-293. PMID 35306217
68. Giralt S, Garderet L, Durie B, et al. American Society of Blood and Marrow Transplantation, European Society of Blood and Marrow Transplantation, Blood and Marrow Transplant Clinical Trials Network, and International Myeloma Working Group Consensus Conference on Salvage Hematopoietic Cell Transplantation in Patients with Relapsed Multiple Myeloma. *Biol Blood Marrow Transplant.* Dec 2015; 21(12): 2039-2051. PMID 26428082
69. Kanate AS, Majhail NS, Savani BN, et al. Indications for Hematopoietic Cell Transplantation and Immune Effector Cell Therapy: Guidelines from the American Society for Transplantation and Cellular Therapy. *Biol Blood Marrow Transplant.* Jul 2020; 26(7): 1247-1256. PMID 32165328
70. Lokhorst H, Einsele H, Vesole D, et al. International Myeloma Working Group consensus statement regarding the current status of allogeneic stem-cell transplantation for multiple myeloma. *J Clin Oncol.* Oct 10 2010; 28(29): 4521-30. PMID 20697091
71. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for Stem Cell Transplantation (Formerly 110.8.1) (110.23). 2016; <https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?NCDId=366&ncdver=1&bc=AAAAIAAAAA&> Accessed November 27, 2023.