



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Oncologic Applications of Photodynamic Therapy, Including Barrett Esophagus

Table of Contents

- [Policy: Commercial](#)
- [Coding Information](#)
- [Information Pertaining to All Policies](#)
- [Policy: Medicare](#)
- [Description](#)
- [References](#)
- [Authorization Information](#)
- [Policy History](#)

Policy Number: 454

BCBSA Reference Number: 8.01.06 (For Plan internal use only)
NCD/LCD: NA

Related Policies

- Dermatologic Applications of Photodynamic Therapy, #[463](#)
- Endoscopic Radiofrequency Ablation or Cryoablation for Treatment of Barrett's Esophagus, #[218](#)
- Photodynamic Therapy for Choroidal Neovascularization, #[600](#)
- Focal Treatments for Prostate Cancer, #[733](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

One or more courses of photodynamic therapy may be considered **MEDICALLY NECESSARY** for the following oncologic applications:

- Palliative treatment of obstructing esophageal cancer
- Palliative treatment of obstructing endobronchial lesions
- Treatment of early-stage non-small cell lung cancer in individuals who are ineligible for surgery and radiation therapy
- Treatment of high-grade dysplasia in Barrett esophagus
- Palliative treatment of unresectable cholangiocarcinoma when used with stenting.

Other oncologic applications of photodynamic therapy are **INVESTIGATIONAL** including, but not limited to, other malignancies and Barrett esophagus without associated high-grade dysplasia.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	Prior authorization is not required .
Commercial PPO and Indemnity	Prior authorization is not required .
Medicare HMO BlueSM	Prior authorization is not required .
Medicare PPO BlueSM	Prior authorization is not required .

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The above **medical necessity criteria MUST** be met for the following codes to be covered for **Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:**

CPT Codes

CPT codes:	Code Description
31641	Bronchoscopy, rigid or flexible, including fluoroscopic guidance, when performed; with destruction of tumor or relief of stenosis by any method other than excision (eg, laser therapy, cryotherapy)
43229	Esophagoscopy, flexible, transoral; with ablation of tumor(s), polyp(s), or other lesion(s) (includes pre- and post-dilation and guide wire passage, when performed)
96570	Photodynamic therapy by endoscopic application of light to ablate abnormal tissue via activation of photosensitive drug(s); first 30 minutes (List separately in addition to code for endoscopy or bronchoscopy)
96571	Photodynamic therapy by endoscopic application of light to ablate abnormal tissue via activation of photosensitive drug(s); each additional 15 minutes

HCPCS Codes

HCPCS codes:	Code Description
J9600	Porfimer sodium, 75 mg

The following ICD Diagnosis Codes are considered medically necessary when submitted with the CPT and HCPCS codes above if **medical necessity criteria** are met:

ICD-10 Diagnosis Codes

ICD-10-CM Diagnosis codes:	Code Description
C15.3	Malignant neoplasm of upper third of esophagus
C15.4	Malignant neoplasm of middle third of esophagus
C15.5	Malignant neoplasm of lower third of esophagus
C15.8	Malignant neoplasm of overlapping sites of esophagus
C15.9	Malignant neoplasm of esophagus, unspecified

C22.1	Intrahepatic bile duct carcinoma
C24.0	Malignant neoplasm of extrahepatic bile duct
C24.1	Malignant neoplasm of ampulla of Vater
C24.8	Malignant neoplasm of overlapping sites of biliary tract
C24.9	Malignant neoplasm of biliary tract, unspecified
C34.00	Malignant neoplasm of unspecified main bronchus
C34.01	Malignant neoplasm of right main bronchus
C34.02	Malignant neoplasm of left main bronchus
C34.10	Malignant neoplasm of upper lobe, unspecified bronchus or lung
C34.11	Malignant neoplasm of upper lobe, right bronchus or lung
C34.12	Malignant neoplasm of upper lobe, left bronchus or lung
C34.2	Malignant neoplasm of middle lobe, bronchus or lung
C34.30	Malignant neoplasm of lower lobe, unspecified bronchus or lung
C34.31	Malignant neoplasm of lower lobe, right bronchus or lung
C34.32	Malignant neoplasm of lower lobe, left bronchus or lung
C34.80	Malignant neoplasm of overlapping sites of unspecified bronchus and lung
C34.81	Malignant neoplasm of overlapping sites of right bronchus and lung
C34.82	Malignant neoplasm of overlapping sites of left bronchus and lung
C34.90	Malignant neoplasm of unspecified part of unspecified bronchus or lung
C34.91	Malignant neoplasm of unspecified part of right bronchus or lung
C34.92	Malignant neoplasm of unspecified part of left bronchus or lung
C78.00	Secondary malignant neoplasm of unspecified lung
C78.01	Secondary malignant neoplasm of right lung
C78.02	Secondary malignant neoplasm of left lung
C78.7	Secondary malignant neoplasm of liver and intrahepatic bile duct
C78.80	Secondary malignant neoplasm of unspecified digestive organ
C78.89	Secondary malignant neoplasm of other digestive organs
D00.1	Carcinoma in situ of esophagus
D02.20	Carcinoma in situ of unspecified bronchus and lung
D02.21	Carcinoma in situ of right bronchus and lung
D02.22	Carcinoma in situ of left bronchus and lung
K22.711	Barrett's esophagus with high grade dysplasia

Description

Photodynamic Therapy

Photodynamic therapy (PDT) has been investigated for use in a wide variety of tumors, including esophageal, lung, cholangiocarcinoma, prostate, bladder, breast, brain (administered intraoperatively), skin, and head and neck cancers. Barrett esophagus also has been treated with PDT. PDT for focal treatment of prostate cancer is discussed in policy #733

Several photosensitizing agents have been used in PDT: porfimer sodium (Photofrin®), administered intravenously 48 hours before light exposure, and 5-aminolevulinic acid, administered orally 4 to 6 hours before the procedure. Aminolevulinic acid is metabolized to protoporphyrin IX, which is preferentially taken up by the mucosa. Clearance of porfimer occurs in a variety of normal tissues over 40 to 72 hours, but tumor cells retain porfimer for a longer period. Laser treatment of Barrett esophagus may be enhanced by the use of balloons containing a cylindrical diffusing fiber. The balloon compresses the mucosal folds of the esophagus, thus increasing the likelihood that the entire Barrett mucosa is exposed to light. All patients who receive porfimer become photosensitive and must avoid exposure of skin and eyes to direct sunlight or bright indoor light for 30 days.

Summary

Description

Photodynamic therapy (PDT; also called phototherapy, photoradiotherapy, photosensitizing therapy, or photochemotherapy) is an ablative treatment that uses a photosensitizing agent to expose tumor cells to a light source of a specific wavelength for the purpose of damaging the cells. After administration of the photosensitizing agent, the target tissue is exposed to light using a variety of laser techniques. For example, a laser fiber may be placed through the channel of the endoscope, or a specialized modified diffuser may be placed via fluoroscopic guidance. Treatment for tumor cells occurs through selective retention of the photosensitizing agent and the selective delivery of light.

Summary of Evidence

For individuals who have obstructing esophageal cancer who receive photodynamic therapy (PDT) as palliation, the evidence includes systematic reviews, randomized controlled trials (RCTs), and uncontrolled single-arm studies. Relevant outcomes are change in disease status, symptoms, quality of life, and treatment-related morbidity. A meta-analysis comparing PDT with neodymium-doped yttrium aluminum garnet (Nd:YAG) laser suggested that improvements in dysphagia are similar, although estimates are imprecise. Compared with the Nd:YAG laser, PDT is associated with a lower risk of perforation and a higher risk of adverse reactions to the light (e.g. photosensitivity). PDT plus argon plasma coagulation appears to prolong the time to recurrence of dysphagia as opposed to argon plasma coagulation alone. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have obstructing endobronchial lesions who receive PDT as palliation, the evidence includes RCTs and uncontrolled single-arm studies. Relevant outcomes are change in disease status, symptoms, quality of life, and treatment-related morbidity. Evidence from RCTs comparing PDT with Nd:YAG laser has generally supported reductions in symptoms using PDT similar to those using a laser. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have early-stage non-small-cell lung cancer (NSCLC) who are not candidates for surgery or radiotherapy who receive PDT, the evidence includes uncontrolled single-arm studies. Relevant outcomes are overall survival (OS), disease-specific survival, change in disease status, quality of life, and treatment-related morbidity. There are few patients with early-stage NSCLC who are not candidates for surgery or radiotherapy. While several treatment methods (eg, laser, electrocautery, cryotherapy, brachytherapy) are available for this population, studies comparing the treatment methods are not available. Case series of PDT include between 21 and 95 patients and have reported complete response rates ranging from 72% to 100%. Given the small size of this potential population and the ineligibility for standard surgical treatment or radiotherapy, it is unlikely that stronger evidence will become available. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with Barrett esophagus with high-grade dysplasia who receive PDT, the evidence includes 2 systematic reviews and 2 RCTs. Relevant outcomes are OS, disease-specific survival, change in disease status, quality of life, and treatment-related morbidity. One RCT compared PDT plus a proton pump inhibitor with a proton pump inhibitor alone and demonstrated higher response rates and lower risk of progression with cancer persisting during 5 years of follow-up for patients in the PDT plus proton inhibitor group. The results of the RCT also revealed that patients treated with PDT had significantly more complications, including a high rate of strictures. Another RCT compared PDT performed with different photosensitizers; results revealed that neither were valuable long-term treatments for dysplastic Barrett esophagus. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome. For individuals who have unresectable cholangiocarcinoma who receive PDT plus stenting as palliation, the evidence includes systematic reviews, RCTs, and observational studies. Relevant outcomes are change in disease status, symptoms, quality of life, and treatment-related morbidity. Three small RCTs and several observational studies have found that PDT plus stenting is associated with the greater elimination of bile duct stenosis and improved survival benefit compared with stenting alone. One RCT comparing stenting plus chemotherapy and PDT with stenting plus chemotherapy without PDT reported longer progression-free survival, but not OS, with similar adverse event rates. Case series have suggested an improvement in the quality of life with PDT. The main complication of PDT in cholangiocarcinoma is cholangitis. Given the small size of this potential population, it is unlikely that stronger evidence will become available. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have other malignancies (eg, gynecologic, bladder, head and neck, brain, soft tissue) who receive PDT, the evidence includes controlled observational studies and uncontrolled single-arm studies. Relevant outcomes are OS, disease-specific survival, change in disease status, quality of life, and treatment-related morbidity. The published literature on PDT for these malignancies is generally comprised of small case series without comparator groups. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Policy History

Date	Action
9/2023	Annual policy review. Reference added. Minor editorial refinements to policy statements; intent unchanged.
9/2022	Annual policy review. Reference added. Minor editorial refinements to policy statements; intent unchanged.
8/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
9/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
9/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
9/2018	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
1/2018	Annual policy review. New medically necessary indications described. Clarified coding information. Effective 1/1/2018.
5/2015	Annual policy review. New references added. Clarified coding language.
7/2014	Updated Coding section with ICD10 procedure and diagnosis codes. Effective 10/2015.
5/2014	Annual policy review. New references added.
1/2014	Added new CPT code 43229 and removed deleted code 43228.
6/2013	Annual policy review. New references added.
5/2013	Annual policy review. New references added.
11/2011-4/2012	Medical policy ICD10 remediation: Formatting, editing and coding updates. No changes to policy statements.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Pinnacle Biologics. Photofrin (porfimer sodium) Injection [prescribing information]. 2019; https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/020451s029,021525s005lbl.pdf. Accessed May 23, 2023.
2. Fayer D, Corbett M, Heirs M, et al. A systematic review of photodynamic therapy in the treatment of pre-cancerous skin conditions, Barrett's oesophagus and cancers of the biliary tract, brain, head and neck, lung, oesophagus and skin. *Health Technol Assess*. Jul 2010; 14(37): 1-288. PMID 20663420
3. Dai Y, Li C, Xie Y, et al. Interventions for dysphagia in oesophageal cancer. *Cochrane Database Syst Rev*. Oct 30 2014; 2014(10): CD005048. PMID 25354795
4. Lightdale CJ, Heier SK, Marcon NE, et al. Photodynamic therapy with porfimer sodium versus thermal ablation therapy with Nd:YAG laser for palliation of esophageal cancer: a multicenter randomized trial. *Gastrointest Endosc*. Dec 1995; 42(6): 507-12. PMID 8674919

5. Heier SK, Rothman KA, Heier LM, et al. Photodynamic therapy for obstructing esophageal cancer: light dosimetry and randomized comparison with Nd:YAG laser therapy. *Gastroenterology*. Jul 1995; 109(1): 63-72. PMID 7541003
6. Rupinski M, Zagorowicz E, Regula J, et al. Randomized comparison of three palliative regimens including brachytherapy, photodynamic therapy, and APC in patients with malignant dysphagia (CONSORT 1a) (Revised II). *Am J Gastroenterol*. Sep 2011; 106(9): 1612-20. PMID 21670770
7. McCann P, Stafinski T, Wong C, et al. The safety and effectiveness of endoscopic and non-endoscopic approaches to the management of early esophageal cancer: a systematic review. *Cancer Treat Rev*. Feb 2011; 37(1): 11-62. PMID 20570442
8. Li LB, Xie JM, Zhang XN, et al. Retrospective study of photodynamic therapy vs photodynamic therapy combined with chemotherapy and chemotherapy alone on advanced esophageal cancer. *Photodiagnosis Photodyn Ther*. Sep 2010; 7(3): 139-43. PMID 20728836
9. Akopov A, Rusanov A, Gerasin A, et al. Preoperative endobronchial photodynamic therapy improves resectability in initially irresectable (inoperable) locally advanced non small cell lung cancer. *Photodiagnosis Photodyn Ther*. Sep 2014; 11(3): 259-64. PMID 24704942
10. Diaz-Jiménez JP, Martínez-Ballarín JE, Lluell A, et al. Efficacy and safety of photodynamic therapy versus Nd-YAG laser resection in NSCLC with airway obstruction. *Eur Respir J*. Oct 1999; 14(4): 800-5. PMID 10573224
11. Kato H, Okunaka T, Shimatani H. Photodynamic therapy for early stage bronchogenic carcinoma. *J Clin Laser Med Surg*. Oct 1996; 14(5): 235-8. PMID 9612188
12. Endo C, Miyamoto A, Sakurada A, et al. Results of long-term follow-up of photodynamic therapy for roentgenographically occult bronchogenic squamous cell carcinoma. *Chest*. Aug 2009; 136(2): 369-375. PMID 19318660
13. Moghissi K, Dixon K, Thorpe JA, et al. Photodynamic therapy (PDT) in early central lung cancer: a treatment option for patients ineligible for surgical resection. *Thorax*. May 2007; 62(5): 391-5. PMID 17090572
14. Corti L, Toniolo L, Boso C, et al. Long-term survival of patients treated with photodynamic therapy for carcinoma in situ and early non-small-cell lung carcinoma. *Lasers Surg Med*. Jun 2007; 39(5): 394-402. PMID 17565719
15. Furukawa K, Kato H, Konaka C, et al. Locally recurrent central-type early stage lung cancer 1.0 cm in diameter after complete remission by photodynamic therapy. *Chest*. Nov 2005; 128(5): 3269-75. PMID 16306036
16. Cortese DA, Edell ES, Kinsey JH. Photodynamic therapy for early stage squamous cell carcinoma of the lung. *Mayo Clin Proc*. Jul 1997; 72(7): 595-602. PMID 9212759
17. Spechler SJ, Sharma P, Souza RF, et al. American Gastroenterological Association medical position statement on the management of Barrett's esophagus. *Gastroenterology*. Mar 2011; 140(3): 1084-91. PMID 21376940
18. Konda VJ, Waxman I. Endotherapy for Barrett's esophagus. *Am J Gastroenterol*. Jun 2012; 107(6): 827-33. PMID 22488078
19. Overholt BF, Wang KK, Burdick JS, et al. Five-year efficacy and safety of photodynamic therapy with Photofrin in Barrett's high-grade dysplasia. *Gastrointest Endosc*. Sep 2007; 66(3): 460-8. PMID 17643436
20. Dunn JM, Mackenzie GD, Banks MR, et al. A randomised controlled trial of ALA vs. Photofrin photodynamic therapy for high-grade dysplasia arising in Barrett's oesophagus. *Lasers Med Sci*. May 2013; 28(3): 707-15. PMID 22699800
21. Kohoutova D, Haidry R, Banks M, et al. Long-term outcomes of the randomized controlled trial comparing 5-aminolaevulinic acid and Photofrin photodynamic therapy for Barrett's oesophagus related neoplasia. *Scand J Gastroenterol*. May 2018; 53(5): 527-532. PMID 29161901
22. Gao F, Bai Y, Ma SR, et al. Systematic review: photodynamic therapy for unresectable cholangiocarcinoma. *J Hepatobiliary Pancreat Sci*. Mar 2010; 17(2): 125-31. PMID 19455276
23. Tomizawa Y, Tian J. Photodynamic therapy for unresectable cholangiocarcinoma. *Dig Dis Sci*. Feb 2012; 57(2): 274-83. PMID 22057285
24. Lu Y, Liu L, Wu JC, et al. Efficacy and safety of photodynamic therapy for unresectable cholangiocarcinoma: A meta-analysis. *Clin Res Hepatol Gastroenterol*. Dec 2015; 39(6): 718-24. PMID 26070572
25. Mohan BP, Chandan S, Khan SR, et al. Photodynamic Therapy (PDT), Radiofrequency Ablation (RFA) With Biliary Stents in Palliative Treatment of Unresectable Extrahepatic Cholangiocarcinoma: A

- Systematic Review and Meta-analysis. *J Clin Gastroenterol*. Feb 01 2022; 56(2): e153-e160. PMID 33780214
26. Ortner ME, Caca K, Berr F, et al. Successful photodynamic therapy for nonresectable cholangiocarcinoma: a randomized prospective study. *Gastroenterology*. Nov 2003; 125(5): 1355-63. PMID 14598251
 27. Zoepf T, Jakobs R, Arnold JC, et al. Palliation of nonresectable bile duct cancer: improved survival after photodynamic therapy. *Am J Gastroenterol*. Nov 2005; 100(11): 2426-30. PMID 16279895
 28. Hauge T, Hauge PW, Warloe T, et al. Randomised controlled trial of temoporfin photodynamic therapy plus chemotherapy in nonresectable biliary carcinoma--PCS Nordic study. *Photodiagnosis Photodyn Ther*. Mar 2016; 13: 330-333. PMID 26415549
 29. Pereira SP, Aithal GP, Ragunath K, et al. Safety and long term efficacy of porfimer sodium photodynamic therapy in locally advanced biliary tract carcinoma. *Photodiagnosis Photodyn Ther*. Dec 2012; 9(4): 287-92. PMID 23200007
 30. Shim CS, Cheon YK, Cha SW, et al. Prospective study of the effectiveness of percutaneous transhepatic photodynamic therapy for advanced bile duct cancer and the role of intraductal ultrasonography in response assessment. *Endoscopy*. May 2005; 37(5): 425-33. PMID 15844020
 31. Harewood GC, Baron TH, Rumalla A, et al. Pilot study to assess patient outcomes following endoscopic application of photodynamic therapy for advanced cholangiocarcinoma. *J Gastroenterol Hepatol*. Mar 2005; 20(3): 415-20. PMID 15740486
 32. Berr F. Photodynamic therapy for cholangiocarcinoma. *Semin Liver Dis*. May 2004; 24(2): 177-87. PMID 15192790
 33. Baron TH. Photodynamic therapy: standard of care for palliation of cholangiocarcinoma?. *Clin Gastroenterol Hepatol*. Mar 2008; 6(3): 266-7. PMID 18328433
 34. Godoy H, Vaddadi P, Cooper M, et al. Photodynamic therapy effectively palliates gynecologic malignancies. *Eur J Gynaecol Oncol*. 2013; 34(4): 300-2. PMID 24020133
 35. Choi MC, Jung SG, Park H, et al. Fertility preservation via photodynamic therapy in young patients with early-stage uterine endometrial cancer: a long-term follow-up study. *Int J Gynecol Cancer*. May 2013; 23(4): 698-704. PMID 23478222
 36. Choi MC, Jung SG, Park H, et al. Fertility preservation by photodynamic therapy combined with conization in young patients with early stage cervical cancer: a pilot study. *Photodiagnosis Photodyn Ther*. Sep 2014; 11(3): 420-5. PMID 24927981
 37. Zhang W, Zhang A, Sun W, et al. Efficacy and safety of photodynamic therapy for cervical intraepithelial neoplasia and human papilloma virus infection: A systematic review and meta-analysis of randomized clinical trials. *Medicine (Baltimore)*. May 2018; 97(21): e10864. PMID 29794788
 38. Tao XH, Guan Y, Shao D, et al. Efficacy and safety of photodynamic therapy for cervical intraepithelial neoplasia: a systemic review. *Photodiagnosis Photodyn Ther*. Jun 2014; 11(2): 104-12. PMID 24631593
 39. Hillemanns P, Garcia F, Petry KU, et al. A randomized study of hexaminolevulinate photodynamic therapy in patients with cervical intraepithelial neoplasia 1/2. *Am J Obstet Gynecol*. Apr 2015; 212(4): 465.e1-7. PMID 25467012
 40. Istomin YP, Lapzevich TP, Chalau VN, et al. Photodynamic therapy of cervical intraepithelial neoplasia grades II and III with Photolon. *Photodiagnosis Photodyn Ther*. Sep 2010; 7(3): 144-51. PMID 20728837
 41. Soergel P, Dahl GF, Onsrud M, et al. Photodynamic therapy of cervical intraepithelial neoplasia 1-3 and human papilloma virus (HMPV) infection with methylaminolevulinate and hexaminolevulinate--a double-blind, dose-finding study. *Lasers Surg Med*. Aug 2012; 44(6): 468-74. PMID 22693121
 42. Winters U, Daayana S, Lear JT, et al. Clinical and immunologic results of a phase II trial of sequential imiquimod and photodynamic therapy for vulval intraepithelial neoplasia. *Clin Cancer Res*. Aug 15 2008; 14(16): 5292-9. PMID 18698049
 43. Zhang R, Wang L. Photodynamic therapy for treatment of usual-type vulvar intraepithelial neoplasia: a case report and literature review. *J Int Med Res*. Aug 2019; 47(8): 4019-4026. PMID 31364444
 44. Bader MJ, Stepp H, Beyer W, et al. Photodynamic therapy of bladder cancer - a phase I study using hexaminolevulinate (HAL). *Urol Oncol*. Oct 2013; 31(7): 1178-83. PMID 22440147
 45. Lee JY, Diaz RR, Cho KS, et al. Efficacy and safety of photodynamic therapy for recurrent, high grade nonmuscle invasive bladder cancer refractory or intolerant to bacille Calmette-Guérin immunotherapy. *J Urol*. Oct 2013; 190(4): 1192-9. PMID 23648222

46. Gondivkar SM, Gadbail AR, Choudhary MG, et al. Photodynamic treatment outcomes of potentially-malignant lesions and malignancies of the head and neck region: A systematic review. *J Investig Clin Dent*. Feb 2018; 9(1). PMID 28480637
47. de Visscher SA, Dijkstra PU, Tan IB, et al. mTHPC mediated photodynamic therapy (PDT) of squamous cell carcinoma in the head and neck: a systematic review. *Oral Oncol*. Mar 2013; 49(3): 192-210. PMID 23068024
48. Wildeman MA, Nyst HJ, Karakullukcu B, et al. Photodynamic therapy in the therapy for recurrent/persistent nasopharyngeal cancer. *Head Neck Oncol*. Dec 17 2009; 1: 40. PMID 20017928
49. Karakullukcu B, Stoker SD, Wildeman AP, et al. A matched cohort comparison of mTHPC-mediated photodynamic therapy and trans-oral surgery of early stage oral cavity squamous cell cancer. *Eur Arch Otorhinolaryngol*. Mar 2013; 270(3): 1093-7. PMID 22773192
50. Ahn PH, Quon H, O'Malley BW, et al. Toxicities and early outcomes in a phase 1 trial of photodynamic therapy for premalignant and early stage head and neck tumors. *Oral Oncol*. Apr 2016; 55: 37-42. PMID 26865261
51. Biel MA. Photodynamic therapy treatment of early oral and laryngeal cancers. *Photochem Photobiol*. 2007; 83(5): 1063-8. PMID 17880501
52. Silbergleit AK, Somers ML, Schweitzer VG, et al. Vocal fold vibration after photofrin-mediated photodynamic therapy for treatment of early-stage laryngeal malignancies. *J Voice*. Nov 2013; 27(6): 762-4. PMID 24119638
53. Wildeman MA, Fles R, Herdini C, et al. Primary treatment results of Nasopharyngeal Carcinoma (NPC) in Yogyakarta, Indonesia. *PLoS One*. 2013; 8(5): e63706. PMID 23675501
54. Durbec M, Cosmidis A, Fuchsmann C, et al. Efficacy and safety of photodynamic therapy with temoporfin in curative treatment of recurrent carcinoma of the oral cavity and oropharynx. *Eur Arch Otorhinolaryngol*. Mar 2013; 270(4): 1433-9. PMID 22927020
55. Rigual NR, Shafirstein G, Frustino J, et al. Adjuvant intraoperative photodynamic therapy in head and neck cancer. *JAMA Otolaryngol Head Neck Surg*. Jul 2013; 139(7): 706-11. PMID 23868427
56. Rigual NR, Thankappan K, Cooper M, et al. Photodynamic therapy for head and neck dysplasia and cancer. *Arch Otolaryngol Head Neck Surg*. Aug 2009; 135(8): 784-8. PMID 19687399
57. Schweitzer VG, Somers ML. PHOTOFRIN-mediated photodynamic therapy for treatment of early stage (Tis-T2N0M0) SqCCa of oral cavity and oropharynx. *Lasers Surg Med*. Jan 2010; 42(1): 1-8. PMID 20077493
58. Lambert A, Nees L, Nuyts S, et al. Photodynamic Therapy as an Alternative Therapeutic Tool in Functionally Inoperable Oral and Oropharyngeal Carcinoma: A Single Tertiary Center Retrospective Cohort Analysis. *Front Oncol*. 2021; 11: 626394. PMID 33747943
59. Muragaki Y, Akimoto J, Maruyama T, et al. Phase II clinical study on intraoperative photodynamic therapy with talaporfin sodium and semiconductor laser in patients with malignant brain tumors. *J Neurosurg*. Oct 2013; 119(4): 845-52. PMID 23952800
60. Aziz F, Telara S, Moseley H, et al. Photodynamic therapy adjuvant to surgery in metastatic carcinoma in brain. *Photodiagnosis Photodyn Ther*. 2009; 6(3-4): 227-30. PMID 19932456
61. Nakamura T, Kusuzaki K, Matsubara T, et al. Long-term clinical outcome in patients with high-grade soft tissue sarcoma who were treated with surgical adjuvant therapy using acridine orange after intra-lesional or marginal resection. *Photodiagnosis Photodyn Ther*. Sep 2018; 23: 165-170. PMID 29885811
62. FNCLCC. The Free Dictionary by Farlex. <https://acronyms.thefreedictionary.com/FNCLCC>. Accessed May 23, 2023.
63. Matsubara T, Kusuzaki K, Matsumine A, et al. Can a less radical surgery using photodynamic therapy with acridine orange be equal to a wide-margin resection?. *Clin Orthop Relat Res*. Mar 2013; 471(3): 792-802. PMID 23008027
64. Matzi V, Maier A, Woltsche M, et al. Polyhematoporphyrin-mediated photodynamic therapy and decortication in palliation of malignant pleural mesothelioma: a clinical pilot study. *Interact Cardiovasc Thorac Surg*. Mar 2004; 3(1): 52-6. PMID 17670175
65. Lindenmann J, Matzi V, Neuboeck N, et al. Multimodal therapy of malignant pleural mesothelioma: is the replacement of radical surgery imminent?. *Interact Cardiovasc Thorac Surg*. Mar 2013; 16(3): 237-43. PMID 23171517
66. Friedberg JS, Simone CB, Culligan MJ, et al. Extended Pleurectomy-Decortication-Based Treatment for Advanced Stage Epithelial Mesothelioma Yielding a Median Survival of Nearly Three Years. *Ann Thorac Surg*. Mar 2017; 103(3): 912-919. PMID 27825687

67. Pereira S. Photodynamic therapy for pancreatic and biliary tract cancer: the United Kingdom experience. *J Natl Compr Canc Netw*. Oct 01 2012; 10 Suppl 2: S48-51. PMID 23055216
68. Huggett MT, Jermyn M, Gillams A, et al. Phase I/II study of verteporfin photodynamic therapy in locally advanced pancreatic cancer. *Br J Cancer*. Apr 02 2014; 110(7): 1698-704. PMID 24569464
69. Bahng S, Yoo BC, Paik SW, et al. Photodynamic therapy for bile duct invasion of hepatocellular carcinoma. *Photochem Photobiol Sci*. Mar 2013; 12(3): 439-45. PMID 23175171
70. Vohra F, Al-Kheraif AA, Qadri T, et al. Efficacy of photodynamic therapy in the management of oral premalignant lesions. A systematic review. *Photodiagnosis Photodyn Ther*. Mar 2015; 12(1): 150-9. PMID 25315968
71. Wisnivesky JP, Yung RC, Mathur PN, et al. Diagnosis and treatment of bronchial intraepithelial neoplasia and early lung cancer of the central airways: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest*. May 2013; 143(5 Suppl): e263S-e277S. PMID 23649442
72. Shaheen NJ, Falk GW, Iyer PG, et al. ACG Clinical Guideline: Diagnosis and Management of Barrett's Esophagus. *Am J Gastroenterol*. Jan 2016; 111(1): 30-50; quiz 51. PMID 26526079
73. Shaheen NJ, Falk GW, Iyer PG, et al. Diagnosis and Management of Barrett's Esophagus: An Updated ACG Guideline. *Am J Gastroenterol*. Apr 01 2022; 117(4): 559-587. PMID 35354777
74. Sharma P, Shaheen NJ, Katzka D, et al. AGA Clinical Practice Update on Endoscopic Treatment of Barrett's Esophagus With Dysplasia and/or Early Cancer: Expert Review. *Gastroenterology*. Feb 2020; 158(3): 760-769. PMID 31730766
75. Ahmed O, Lee JH, Thompson CC, et al. AGA Clinical Practice Update on the Optimal Management of the Malignant Alimentary Tract Obstruction: Expert Review. *Clin Gastroenterol Hepatol*. Sep 2021; 19(9): 1780-1788. PMID 33813072
76. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Esophageal and esophagogastric junction cancer. Version 2.2023. https://www.nccn.org/professionals/physician_gls/pdf/esophageal.pdf. Accessed May 23, 2023.
77. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Biliary tract cancers. Version 2.2023. https://www.nccn.org/professionals/physician_gls/pdf/btc.pdf. Accessed May 22, 2023.
78. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Non-small cell lung cancer. Version 3.2023. https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf. Accessed May 21, 2023.
79. National Institute for Health and Care Excellence. Photodynamic therapy for bile duct cancer [IPG134]. 2005; <https://www.nice.org.uk/guidance/ipg134>. Accessed May 19, 2023.
80. National Institute for Health and Care Excellence. Photodynamic therapy for localised inoperable endobronchial cancer [IPG137]. 2005; <http://www.nice.org.uk/guidance/ipg137>. Accessed May 16, 2023.
81. National Institute for Health and Care Excellence. Photodynamic therapy for advanced bronchial carcinoma [IPG87]. 2004; <https://www.nice.org.uk/guidance/ipg87>. Accessed May 20, 2023.
82. National Institute for Health and Care Excellence. Interstitial photodynamic therapy for malignant parotid tumours [IPG259]. 2008; <http://www.nice.org.uk/nicemedia/pdf/IPG259Guidance.pdf>. Accessed May 23, 2023.
83. National Institute for Health and Care Excellence. Photodynamic therapy for early-stage oesophageal cancer [IPG200]. 2006; <http://www.nice.org.uk/nicemedia/pdf/IPG200guidance.pdf>. Accessed May 17, 2023.
84. National Institute for Health and Care Excellence. Photodynamic therapy for Barrett's oesophagus [IPG350]. 2010; <http://www.nice.org.uk/guidance/ipg350>. Accessed May 21, 2022.
85. National Institute for Health and Care Excellence. Photodynamic therapy for brain tumours [IPG290]. 2009; <http://www.nice.org.uk/nicemedia/pdf/IPG290Guidance.pdf>. Accessed May 18, 2023.