



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an independent
Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Patient-Controlled End of Range Motion Stretching Devices

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 721

BCBSA Reference Number: 1.03.05 (For Plan internal use only)

NCD/LCD: N/A

Related Policies

None

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

Patient-controlled end range of motion stretching devices (static progressive and serial) are considered **INVESTIGATIONAL**.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.
Medicare HMO Blue SM	This is not a covered service.
Medicare PPO Blue SM	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The following HCPCS codes are considered investigational for **Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:**

HCPCS Codes

HCPCS codes:	Code Description
E1801	Static progressive stretch elbow device, extension and/or flexion, with or without range of motion adjustment, includes all components and accessories
E1806	Static progressive stretch wrist device, flexion and/or extension, with or without range of motion adjustment, includes all components and accessories
E1811	Static progressive stretch knee device, extension and/or flexion, with or without range of motion adjustment, includes all components and accessories
E1816	Static progressive stretch ankle device, flexion and/or extension, with or without range of motion adjustment, includes all components and accessories
E1818	Static progressive stretch forearm pronation/supination device, with or without range of motion adjustment, includes all components and accessories
E1831	Static progressive stretch toe device, extension and/or flexion, with or without range of motion adjustment, includes all components and accessories
E1841	Static progressive stretch shoulder device, with or without range of motion adjustment, includes all components and accessories.

Description

Range of Motion Impairments

Loss of full range of motion occurs in a significant proportion of patients following surgical procedures around a joint, such as total knee arthroplasty or anterior cruciate ligament reconstruction. The most common cause of severe postoperative motion loss is the development of intra-articular or extra-articular arthrofibrosis. Arthrofibrosis, characterized by periarticular fibrosis and bands of scar tissue, is described as a painful loss of end range of motion compared with the normal contralateral side. Loss of knee range of motion can lead to impairments in walking, sitting, rising from a chair, and navigating stairs. In 2010, Stephenson et al estimated that based on the annual rates of total knee arthroplasty and anterior cruciate ligament reconstruction, the number of major knee surgery patients affected by arthrofibrosis in the United States would be at least 85,000 per year, and approximately 21,000 patients each year would be at risk of requiring additional surgery.¹

Treatment

Treatment of arthrofibrosis may include physical therapy, manipulation under anesthesia, arthroscopic or open lysis of adhesions, or revision surgery. Conservative treatment typically consists of postoperative physical therapy with pressure stretching techniques and home exercises. When rehabilitation has failed, serial casting, static braces, or dynamic splints that provide low-load prolonged stretch may be used. Dynamic splints use spring loading or elastic bands to provide low-intensity tension (less than that exerted by a physical therapist) and are designed to be worn over relatively long periods (ie, 6 to 8 hours or overnight). The efficacy of a stretching regimen to permanently remodel tissue is considered to be a function of the intensity, length of the session, number of sessions per day, and number of days per week that stretching is performed.²

This evidence review focuses on patient-controlled mechanical devices that provide either moderate- to high-intensity stretch or static progressive stretch in the home. Patient-controlled stretching devices are used at home to increase range of motion in patients who have impaired functional status due to decreased range of motion. We address 2 types of commercially available devices. Static progressive stretch devices (eg, Joint Active Systems [JAS], Static-Pro) provide low- to moderate-intensity stretching with a crank or

ratchet that progressively increases the stretch within each session, and serial stretch devices (eg, End Range of Motion Improvement [ERMI]) use hydraulics to alternate between periods of higher intensity stretch and relaxation.

Improvement in functional outcomes, such as the ability to perform activities of daily living, is the primary goal of this intervention. Joint range of motion is an intermediate outcome. In 2000, 1 small study by Rowe et al. correlated knee range of motion with functional parameters and concluded that 110° is considered the functional range of motion necessary to allow patients to perform common activities of daily living such as navigating stairs, rising from a low chair or commode, entering or exiting a car, or tying one's shoes.³ This threshold of range of motion is therefore used as a measure of treatment success for individual patients. Loss of knee range of motion of more than 15°, which occurs in about 1% to 2% of patients after anterior cruciate ligament reconstruction, has been associated with loss of quadriceps muscle strength and the development of osteoarthritis.⁴ According to the knee examination form developed by the International Knee Documentation Committee (2000), an extension deficit of 6° to 10° or a flexion deficit of 16° to 25° when compared with the noninvolved knee is categorized "abnormal," and an extension deficit of more than 10° or a flexion deficit of more than 25° when compared with the noninvolved knee is categorized "severely abnormal."⁵ Range of motion thresholds in joints other than the knee have been less clearly defined.

Summary

Patient-controlled stretching devices are used at home to increase range of motion in patients who have impaired functional status due to decreased range of motion. We address 2 types of commercially available devices. Static progressive stretch devices (eg, Joint Active Systems, Static-Pro) provide low- to moderate-intensity stretching with a crank or ratchet that progressively increases the stretch within each session, and serial stretch devices (eg, End Range of Motion Improvement [ERMI]) use hydraulics to alternate between periods of higher intensity stretch and relaxation.

Summary of Evidence

For individuals who have functional limitations in range of motion who receive static progressive stretch devices and physical therapy, the evidence includes RCTs, a systematic review, and case series. Relevant outcomes include symptoms, change in disease status, functional outcomes, and quality of life. Three RCTs have evaluated static progressive stretch devices but comparators in each differed (physical therapy, a dynamic splint, and a serial stretch device). The evidence on static progressive stretch devices does not currently support an improvement in pain and function with static progressive stretch compared to alternative treatments. One RCT found greater improvements in range of motion and Western Ontario and McMaster Universities Arthritis Index(WOMAC) scores with serial stretch devices for the knee compared with static progressive stretch devices. Another RCT evaluating static progressive stretch for shoulder adhesive capsulitis found significant differences in shoulder range of motion compared with physical therapy alone at the end of 4 weeks of treatment, with no difference in pain and function. A third RCT found comparable improvements in most outcomes for the static progressive stretch device compared with dynamic splinting, and a systematic review of case reports and series found similar clinical efficacy for increasing elbow range of motion between static progressive stretch devices and dynamic splints. Dynamic splints are used for 8 to 24 hours per day while static progressive stretch devices require several 30 minute sessions. It is not known whether patient compliance is higher with static progressive stretch devices. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have functional limitations in range of motion who receive serial stretch devices and physical therapy, the evidence includes an RCT and observational studies. Relevant outcomes include symptoms, change in disease status, functional outcomes, and quality of life. The best evidence consists of serial stretching with ERMI devices used to treat knee range of motion. One small RCT and a larger retrospective comparative study have reported that high-intensity stretching with ERMI devices improved range of motion more than lower intensity stretching devices in patients who were post-injury or surgery. Other available data consist of retrospective case series that have demonstrated improved range of motion in patients whose range had plateaued with physical therapy. The clinical significance of gains in this surrogate outcome measure is unclear. Further high-quality comparative trials are needed to determine whether these patient-controlled devices improve functional outcomes better than alternative

treatments and identify the patient populations that might benefit. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Policy History

Date	Action
5/2024	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
5/2023	Annual policy review. Minor editorial refinement to policy statement; intent unchanged.
4/2022	Annual policy review. Summary updated. Policy statements unchanged.
4/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
5/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
4/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
5/2017	Annual policy review. Summary section clarified.
9/2016	Annual policy review. Policy title changed to "Patient-Controlled End Range of Motion Stretching Devices." References added.
9/2015	New medical policy describing investigational indications. Effective 9/1/2015.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

- Stephenson JJ, Quimbo RA, Gu T. Knee-attributable medical costs and risk of re-surgery among patients utilizing non-surgical treatment options for knee arthrofibrosis in a managed care population. *Curr Med Res Opin.* May 2010; 26(5): 1109-18. PMID 20225995
- Jacobs CA, Sciascia AD. Factors that influence the efficacy of stretching programs for patients with hypomobility. *Sports Health.* Nov 2011; 3(6): 520-3. PMID 23016052
- Rowe PJ, Myles CM, Walker C, et al. Knee joint kinematics in gait and other functional activities measured using flexible electrogoniometry: how much knee motion is sufficient for normal daily life?. *Gait Posture.* Oct 2000; 12(2): 143-55. PMID 10998612
- Shelbourne KD, Gray T. Minimum 10-year results after anterior cruciate ligament reconstruction: how the loss of normal knee motion compounds other factors related to the development of osteoarthritis after surgery. *Am J Sports Med.* Mar 2009; 37(3): 471-80. PMID 19059893
- International Knee Documentation Committee (IKDC). IKDC: Knee Form. 2000. Accessed January 24, 2024.
- Sodhi N, Yao B, Anis HK, et al. Patient satisfaction and outcomes of static progressive stretch bracing: a 10-year prospective analysis. *Ann Transl Med.* Feb 2019; 7(4): 67. PMID 30963062
- Papotto BA, Mills T. Treatment of severe flexion deficits following total knee arthroplasty: a randomized clinical trial. *Orthop Nurs.* 2012; 31(1): 29-34. PMID 22278649
- Ibrahim MI, Johnson AJ, Pivec R, et al. Treatment of adhesive capsulitis of the shoulder with a static progressive stretch device: a prospective, randomized study. *J Long Term Eff Med Implants.* 2012; 22(4): 281-91. PMID 23662659
- Ibrahim M, Donatelli R, Hellman M, et al. Efficacy of a static progressive stretch device as an adjunct to physical therapy in treating adhesive capsulitis of the shoulder: a prospective, randomised study. *Physiotherapy.* Sep 2014; 100(3): 228-34. PMID 24211154

10. Hussein AZ, Ibrahim MI, Hellman MA, et al. Static progressive stretch is effective in treating shoulder adhesive capsulitis: Prospective, randomized, controlled study with a two-year follow-up. *Eur J Physiother.* Jun 24 2015;17(3):138-147.
11. Lindenhovius AL, Doornberg JN, Brouwer KM, et al. A prospective randomized controlled trial of dynamic versus static progressive elbow splinting for posttraumatic elbow stiffness. *J Bone Joint Surg Am.* Apr 18 2012; 94(8): 694-700. PMID 22517385
12. Bonutti PM, McGrath MS, Ulrich SD, et al. Static progressive stretch for the treatment of knee stiffness. *Knee.* Aug 2008; 15(4): 272-6. PMID 18538574
13. Ulrich SD, Bonutti PM, Seyler TM, et al. Restoring range of motion via stress relaxation and static progressive stretch in posttraumatic elbow contractures. *J Shoulder Elbow Surg.* Mar 2010; 19(2): 196-201. PMID 19959379
14. Müller AM, Sadoghi P, Lucas R, et al. Effectiveness of bracing in the treatment of nonosseous restriction of elbow mobility: a systematic review and meta-analysis of 13 studies. *J Shoulder Elbow Surg.* Aug 2013; 22(8): 1146-52. PMID 23796383
15. McGrath MS, Ulrich SD, Bonutti PM, et al. Static progressive splinting for restoration of rotational motion of the forearm. *J Hand Ther.* 2009; 22(1): 3-8; quiz 9,. PMID 18950990
16. McGrath MS, Ulrich SD, Bonutti PM, et al. Evaluation of static progressive stretch for the treatment of wrist stiffness. *J Hand Surg Am.* Nov 2008; 33(9): 1498-504. PMID 18984330
17. Lucado AM, Li Z, Russell GB, et al. Changes in impairment and function after static progressive splinting for stiffness after distal radius fracture. *J Hand Ther.* 2008; 21(4): 319-25. PMID 19006757
18. Branch TP, Karsch RE, Mills TJ, et al. Mechanical therapy for loss of knee flexion. *Am J Orthop (Belle Mead NJ).* Apr 2003; 32(4): 195-200. PMID 12723771
19. Dempsey AL, Mills T, Karsch RM, et al. Maximizing total end range time is safe and effective for the conservative treatment of frozen shoulder patients. *Am J Phys Med Rehabil.* Sep 2011; 90(9): 738-45. PMID 21430510